

High Resolution CO Imaging of the Molecular Disk around the Jets in KJ Pn 8

T. Forveille

Observatoire de Grenoble, B.P. 53X, 38041 Grenoble Cedex, France

P.J. Huggins

Physics Department, New York University, 4 Washington Place, New York, NY 10003

R. Bachiller

IGN Observatorio Astronómico Nacional, Apt 1143, 28800 Alcalá de Henares, Spain

and

P. Cox

Institut d'Astrophysique Spatiale, Université de Paris XI, 91405 Orsay, France

Institut d'Astrophysique de Paris, 92 bis, Boulevard Arago, 75014 Paris, France

Received _____; accepted _____

ABSTRACT

We report high resolution ($2.5''$) CO $J=1-0$ line imaging which confirms the presence of a molecular disk around the origin of the spectacular, $14' \times 4'$, episodic jets in the planetary nebula KJPN 8. The disk is $30''$ in diameter with an expansion velocity of $\approx 7 \text{ km s}^{-1}$. The axis of the disk is aligned with the youngest and fastest ($\approx 300 \text{ km s}^{-1}$) of the bipolar jets, and there is evidence for interaction between the jets and the disk material. The inner $4''$ of the disk are photo-ionized by the central star. The disk-jet system dominates the environment of this young nebula, and should govern the morphology of KJPN 8 as it evolves to become fully ionized.

Subject headings: circumstellar matter — ISM: jets and outflows — planetary nebulae: individual (KJPN 8) — stars: mass-loss

1. Introduction

The presence of jets in planetary nebulae (PNe) has only recently been widely recognized, but they prove to be of great interest both for the study of jet phenomena and for their effects on PN structure. In PNe with jets, the action of the collimated flows will modify, and in some cases may completely change the traditional picture of PN formation by the two wind model, in which the shaping is produced by the action of an isotropic wind (e.g., Kahn & West 1985, Balick et al. 1987).

Evidence for jets in PNe is found in objects at all stages of evolution. It includes direct observation of elongated, high velocity structures, which may be episodic and exhibit changes in direction (López 1997), as well as other features in the nebulae such as point symmetries (Schwarz et al. 1992) and poly-polar structures (Manchado et al. 1996), which are likely produced by jets or related processes. Observations suggest that jet activity begins at the earliest stages of PN formation, where it will have most impact on the structure of the neutral circumstellar envelope that evolves into the ionized nebula (Huggins et al. 1996). Direct evidence for interaction of the jets with the envelope is seen in high velocity molecular outflows in several proto-PNe and young PNe, the best studied case being CRL 618 (Cernicharo et al. 1989, Neri et al. 1992, Hajian et al. 1996). The extent of the jets in these young objects is, however, usually quite small and not well resolved, so the recent discovery of very extended jets in KJpn 8 provides a unique opportunity to study their effects.

KJpn 8 (PN G112.5–00.1, K3–89) was first identified as a PN by Kazarian & Parsamian (1971) and independently by Kohoutek (1972). The ionized nebula is small, with an angular diameter of $\approx 4''$, but surrounding this López et al. (1995) have discovered an enormous ($\approx 14' \times 4'$), low surface-brightness, bipolar structure. It consists of symmetric jet features with paired emission regions, characteristic of episodic ejection (see Fig. 1).

Optical line ratios of the emission indicate shock excitation, and the high velocities of the jets ($\gtrsim 200 \text{ km s}^{-1}$) have been directly confirmed by radial velocity measurements (López et al. 1997). Proper motions of the jets have also been detected by Meaburn (1997), and lead to a distance estimate for KJPN 8 of 1.6 kpc. At this distance, the extended bipolar structure is $6.5 \times 1.9 \text{ pc}$ in size.

We have recently reported the detection of millimeter CO emission towards KJPN 8, which reveals the presence of molecular gas near the origin of the jets (Huggins et al. 1997). The distribution of the gas was only partially resolved by our $12''$ observations, but suggested a disk morphology. In view of the potential importance of mutual disk-jet interactions for understanding PN formation, we have re-observed the envelope at much higher angular resolution in CO, and in this *Letter* we report the results.

2. Observations

The observations were made in the 115 GHz (2.6 mm) CO $J = 1 - 0$ line during April and May 1997 using the IRAM interferometer (Guilloteau et al. 1992) located at Plateau de Bure, France. The array consisted of five 15 m antennas, equipped with SIS heterodyne receivers. The observations, centered on KJPN 8, were made with two configurations of the array, C2 and D. At the frequency of the CO line the primary beam is $44''$ (FWHM) and the effective velocity resolution of the line observations was 1.6 km s^{-1} . Observations of the continuum at 115 GHz were also made, with a bandwidth of 320 MHz.

The RF passband and amplitude were calibrated using NRAO 530 (flux 5.75 Jy) and 3C454.3 (6.4 Jy), and phase calibration was performed every 20 minutes using J0102+5824 (2.6 Jy). Data obtained using the IRAM 30 m telescope were used for the zero spacing observations. The uv data were Fourier transformed and CLEANed, using the Clark

algorithm, and the restored Gaussian clean beam is $2.8'' \times 2.1''$ at a position angle of 70° .

The results of the CO observations are shown in Figs. 2–4. The continuum at 115 GHz was not detected with an upper limit of $\lesssim 2$ mJy, consistent with the weak continuum emission of 0.8 mJy at 3.5 cm measured by López et al. (1995).

3. Results and discussion

3.1. Properties of the disk

Our high resolution CO observations provide striking evidence for the presence of a disk at the center of KJPN 8. The overall distribution of the molecular gas is shown in the velocity integrated CO map in Fig. 2. Note that this figure corresponds to the small square at the center of Fig. 1. The CO emission shows a roughly elliptical morphology, with an overall extent of $\approx 30'' \times 14''$ at a position angle of 36° . There is a distinct $\approx 4''$ hole at the center which corresponds in size and position with the small ionized nebula.

The detailed structure and kinematics of the molecular gas can be seen in the channel maps in Fig. 3. Inspection of the maps shows that the main distribution of CO is symmetric in both position and velocity about a systemic velocity of $V_o = -35 \text{ km s}^{-1}$ (LSR), in close agreement with the velocity of the the ionized nebula measured using optical lines ($V_o = -34 \text{ km s}^{-1}$, López et al. 1997). At this velocity, the CO emission is divided into two separate parts, with essentially no emission at the map center. At more negative (approaching) velocities, the two regions of CO emission join together towards the north west, and at more positive (receding) velocities, they join together towards the south east. This kinematic structure is the signature of an expanding disk, whose near side tilts away from earth to the north west. This is fully consistent with the integrated intensity map in Fig. 2, where the projected disk is seen as an elliptical structure. From the orientation and

dimensions of the ellipse in Fig. 2, the axis of the disk lies at a position angle of 126° (i.e., along the minor axis of the ellipse) and is tilted away from the line of sight to the north west by $\approx 65^\circ$.

The CO channel map at the systemic velocity of $V_o = -35 \text{ km s}^{-1}$ corresponds to a cross section of the disk in the plane of the sky. The disk is not flat. At the inner radius it is $\approx 4''$ thick, comparable in size to the ionized nebula, but farther out it thickens to $\approx 20''$. Along the disk axis this structure corresponds to bipolar cavities on each sides of the center, with roughly conical shapes and opening angles of 60° – 90° .

Additional perspectives on the disk structure are shown in the velocity strip maps in Fig. 4. The strips through the center along the principal axes (Figs. 4a and 4b) show the central cavity which corresponds to the position of the ionized nebula. The strip along the direction of the disk axis (4b) also clearly shows the tilt of the disk system; the difference in velocity between the two sides gives an expansion velocity for the bulk of the gas of $\approx 7 \text{ km s}^{-1}$ (corrected for inclination), but much higher velocities (up to 28 km s^{-1}) are present near the center, where the surfaces of the disk are seen almost directly along the line of sight. Irregularities in the disk are also seen in the data, the most remarkable being hollow structures within the disk, shown in the strip in Fig. 4c.

The CO emission can also be used to estimate the mass of the disk. For optically thin, LTE emission, the peak 1–0 intensity of 55 K km s^{-1} corresponds to a CO column density of $8 \times 10^{16} \text{ cm}^{-2}$ for an excitation temperature of 25 K which is typical of that found in PNe (Bachiller et al. 1993). The total flux in the line summed over the disk is $9,600 \text{ K km s}^{-1} \text{ arcsec}^2$. Using a representative abundance of $\text{CO}/\text{H} = 3 \times 10^{-4}$ and a distance $d = 1.6 \text{ kpc}$, this corresponds to a mass of $M_m \gtrsim 0.03 M_\odot$, in good agreement with our previous estimate based on the 2–1 line (Huggins et al. 1997).

The observed flux in the 1–0 line is, in fact, similar to that in the 2–1 line, which

suggests that the CO emission is optically thick. Thus our estimate for M_m is a lower limit. The high opacity also suggests an explanation for the asymmetry of the disk seen in Fig. 2, in which the receding limb is brighter than the approaching limb. If the excitation decreases with distance from the star, as expected, the hotter, inner surface of the receding limb will appear brighter than the outer surface of the approaching limb, as observed.

For comparison with the mass of the disk, the mass of the ionized nebula estimated from the radio continuum is $M_i = 5 \times 10^{-4} M_\odot$ for $d = 1.6$ kpc (Huggins et al. 1997). Thus the ratio M_m/M_i is large ($\gtrsim 60$), and the mass is dominated by the mass of the molecular disk.

3.2. Interaction of the disk and the jets

Our observations of the disk in KJ Pn 8 provide important information on the region where the jets originate. Optical images reveal several different jet axes (see Fig. 1) defined by point symmetric emission regions that are produced by episodic ejections with the rotation or wandering of the jets. The longest jet axis (labeled C by López et al. 1995) lies at a position angle of 71° , and the shortest jet axis, with the most intense emission (axis A), lies at a position angle of 121° .

The orientation of the molecular disk that we measure (Sect. 3.1) is indicated in Fig. 1, and clearly corresponds with jet axis A. In addition, according to López et al. (1997), axis A is tilted by 60° away from the line of sight to the north west, in good agreement with our estimate for the tilt of the molecular disk axis. We therefore conclude that jet A and the disk axis are aligned in space, and are physically related. The expansion time scales, $\tau = R/V$, where R is the characteristic size and V is the expansion velocity, are consistent with this. For the A jets, $\tau_j \approx 2,700$ yr (López et al. 1997); for the (half-size) of

the molecular disk $\tau_d \approx 8,000$ yr; and for ionized inner disk, $\tau_i \lesssim 2,000$ yr.

In optical images of KJPN 8 (López et al. 1995), features which connect the heads of the A jets to the center indicate quite a large opening angle $\approx 45^\circ$ – 60° , which fits in well with the large opening angle of the bipolar cavities formed by the molecular disk. Interaction with the disk is signaled by the high velocity molecular gas (the extreme channels in Fig. 3) which is seen at the rims of the disk, and is likely entrained in the sides of the jets. A similar situation probably also exists in the ionized gas closer ($\lesssim 2''$) to the star. The bulk of the ionized gas has a low expansion velocity ($V_e = 5$ – 7.5 km s $^{-1}$, Sabbadin et al. 1986) similar to that of the molecular gas, and represents the photo-ionized inner disk, but high velocity 50–60 km s $^{-1}$ components of [N II] 6584 Å can also be traced into the center (López et al. 1997).

Given the thinness of the disk relative to the extent of the jets, it seems unlikely that the molecular gas is responsible for focusing the jets, as has been suggested in the case of He 3-1475 by Borkowski et al. (1997). A more likely scenario is that a common mechanism induced the mass-loss in the equatorial plane and created the polar jets close to the center – probably in a binary or multiple central star system. Any neutral gas ejected at high latitudes will have been evacuated along the polar axis by the action of the jets, and the lateral pressure in the jet would naturally expand to fill the opening angles of the disk. Consistent with this, part of the structure in the disk appears to be compressed edges or rims. The more irregular structure may result from instabilities, or from variations in the disk ejection process.

The origin of the wandering or precession of jets in PNe is not clear (see, e.g., Livio & Pringle 1997 for a recent model), but our observations provide useful constraints in the case of KJPN 8. From the geometry and kinematics of the disk there is no evidence for alignment or interaction of the molecular gas with the most extended jets along axis C.

These are the oldest jets, with an expansion time scale of $\sim 16,000$ yr, consistent with the idea that they predate the molecular disk. The shape of the envelope of excited gas which surrounds these jets (see, Fig. 2 of López et al. 1995) also shows no trace of the molecular disk orientation, and, if anything, suggests a waist perpendicular to the C axis. This argues against a commonly assumed scenario in which the jets rotate or precess relative to a well defined equatorial plane, and it rather suggests that the jets and their associated equatorial material have been ejected in a series of different directions during the evolution of the system.

Our observations are also relevant to the future evolution of KJPN 8. The observed structure means that the young ionized nebula at the center of the system is currently surrounded by a reservoir of neutral gas in the disk plane and evacuated poles along the axis of the disk. Whether or not the jets continue to be active, the environment of the nebula is largely determined by these structural features, and they are likely to dominate the future morphological evolution of the nebula.

4. Conclusions

Our molecular line imaging reveals in detail the structure and kinematics of the molecular disk around the origin of the remarkable jets in KJPN 8. The axis of the disk is aligned with the youngest of the bipolar jets and there is evidence for disk-jet interaction. The disk and jets are the major structural features around the young nebula, and are likely to dominate the morphology as KJPN 8 evolves further into a fully ionized PN.

We are grateful to the IRAM staff for obtaining the observations, and for their excellent support during the data reduction. This work has been supported in part by NSF grant AST-9617941 (to P.J.H.).

REFERENCES

- Balick, B., Preston, H.L., Icke, V. 1987, AJ, 94, 1641
- Bachiller, R., Huggins, P.J., Cox, P., Forveille, T. 1993, A&A, 267, 177
- Borkowski, K.J., Blondin, J.M., Harrington, J.P. 1997, ApJ, 482, L97
- Cernicharo, J., Guélin, M., Peñalver, J., Martín-Pintado, J, Mauersberger, R. 1989, A&A, 222, L1
- Hajian, A.R., Phillips, J.A., Terzian, Y. 1996, ApJ, 467, 341
- Huggins, P.J., Bachiller, R., Cox, P., Forveille, T. 1996, A&A, 315, 284
- Huggins, P.J., Bachiller, R., Cox, P., Forveille, T. 1997, ApJ, 483, L57
- Kahn, F.D., West, K.A. 1985, MNRAS, 212, 837
- Kazarian, M.A., Parsamian, E.S.P. 1971, Astron. Tsirk., 602, 6
- Kohoutek, L. 1972, A&A, 16, 291
- Livio, M., Pringle, J.E. 1997, ApJ, 486, 835
- López, J.A. 1997, Planetary Nebulae (IAU symposium 180), ed. H.J. Habing & H.J.G.L.M. Lamers (Dordrecht: Kluwer), in press
- López, J.A., Vasquez, R., Rodriguez, L.F. 1995, ApJ, 455, L63
- López, J.A., Meaburn, J., Bryce, M., Rodriguez, L.F. 1997, ApJ, 475, 705
- Manchado, A., Stanghellini, L., Guerrero, M.A. 1996, ApJ, 466, L95
- Meaburn, J. 1997, MNRAS, in press
- Neri, R., Garcia-Burillo, S., Guélin, M., Cernicharo, J., Guilloteau, S., Lucas, R. 1992, A&A, 262, 544
- Sabbadin, F., Strafella, F., Bianchini, A. 1986, A&AS, 65, 259

Schwarz, H.E., Corradi, R.L.M., Melnick, J. 1992, A&AS, 96, 23

Fig. 1.— Outline of the main optical features of KJpN 8, adapted from the $H\alpha$ image by López et al. (1995). The small square at the center corresponds to the map in Fig. 2. The diagonal lines indicate the orientation of the molecular disk axis (see text)

Fig. 2.— Map of the CO(1–0) emission in KJpN 8 in integrated intensity. The co-ordinates of the map center are 23:24:10.440 60:57:31.00 (J2000.0). The cross indicates the center of the ionized nebula, and the inset shows the beam size. The first solid contour and the contour spacing are 7.5 K km s^{-1} ; the dashed contour is negative with the same spacing

Fig. 3.— Channel maps in the CO(1–0) line. The channel separation is 1.6 km s^{-1} . The lowest contour and the contour intervals are 0.5 K

Fig. 4.— Velocity-strip maps in the CO(1–0) line. a) Perpendicular to the disk axis, through the center. b) Along the disk axis. c) Parallel to the disk axis through offset $3''$, $5''$. The lowest contour and the contour intervals are 0.4 K







